

25th International Conference on
HUMAN METABOLIC HEALTH- DIABETES, OBESITY & METABOLISM
&
2nd World Congress on
NUTRITION AND OBESITY PREVENTION
March 21-22, 2019 Dubai, UAE

Time to optimal glycaemic control and prognostic factors among Type 2 diabetes mellitus patients in public teaching hospitals in Addis Ababa, Ethiopia

Tigist W Leulseged¹ and Birhanu T Ayele²

¹St. Paul's Hospital Millennium Medical College, Ethiopia

²Stellenbosch University, South Africa

Background: Diabetes is a chronic, progressive disease characterized by elevated levels of blood glucose. Poorly managed diabetes leads to serious complications and early death. The prevalence of diabetes has been increasing over the past few decades. Ethiopia is one of African countries with the highest number of people living with diabetes. Studies conducted in Ethiopia and other countries mainly focused on level of glycemic control at one point in time. Studies targeting the time that a patient stayed in a poor glycemic level are lacking.

Objective: To estimate time to first optimal glycemic control and to identify prognostic factors among Type-2 Diabetes Mellitus (T2DM) patients in public teaching hospitals in Addis Ababa, Ethiopia.

Methods: A hospital based retrospective chart review study was conducted from April to July 2018 at diabetes clinic of Addis Ababa's public teaching hospitals among randomly selected sample of 685 charts of T2DM patients who were on follow up from January 1, 2013 to June 30, 2017. Data was collected using pretested data abstraction tool. Data was checked, coded and entered to Epi-Info V.7.2.1.0 and exported to SPSS V.23.0 and STATA V.14.1 for analysis. Descriptive statistics is presented with frequency tables, Kaplan-Meier plots and median survival times. Association was done using Log-rank test and Cox proportional hazard survival model, where hazard ratio, P-value and 95% CI for hazard ratio were used for testing significance and interpretation of results.

Results: Median time to first optimal glycemic control among the study population was 9.5 months. The major factors that affect it are age group (HR=0.635, 95% CI: 0.486-0.831 for 50-59 years, HR=0.558, 95% CI: 0.403-0.771 for 60-69 years and HR=0.495, 95% CI: 0.310-0.790 for ≥70 years), diabetes neuropathy (HR=0.502, 95% CI: 0.375-0.672), more than one complication (HR=0.381, 95% CI: 0.177-0.816), hypertension (HR=0.611, 95% CI: 0.486-0.769), dyslipidemia (HR=0.609, 95% CI: 0.450-0.824), cardiovascular disease (HR=0.670, 95% CI: 0.458-0.979) and hospital patient treated at (HR=1.273, 95% CI: 1.052-1.541).

Conclusion: Median time to first optimal glycemic control among T2DM patients is longer than expected which might imply that patients are being exposed to more risk of complication and death.

Biography

Tigist W. Leulseged is a medical doctor and public health specialist currently working as a lecture at Endocrinology unit under Department of Internal Medicine, St. Paul's Hospital Millennium Medical College in Addis Ababa, Ethiopia. Her research interests are chronic illnesses particularly diabetes epidemiology including gestational diabetes, treatment, patient involvement in effective management and quality of services provided. She has experience in related research areas in both hospitals and teaching institutions.

birtesh@gmail.com
tigidolly@gmail.com

Notes: