

Candidal Onychomycosis in 22 Years Old Saudi Female

Remah Alzayyat*, Abdullah aljaflan and Sarah Albreiki

Department of Dermatology, College of Medicine, ImamAbdulrahman bin Faisal University, Dammam, SaudiArabia

Corresponding Author*

Remah Alzayyat,
Department of Dermatology,
College of Medicine,
Imam Abdulrahman bin Faisal University
Dammam, Saudi Arabia
E-mail: remah.business@gmail.com

Copyright: © 2025 Alzayyat R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: August 08, 2024, Manuscript No. DMCR-24-33332; **Editor assigned:** August 12, 2024, PreQC No. DMCR-24-33332 (PQ); **Reviewed:** August 26, 2024, QC No. DMCR-24-33332; **Revised:** August 03, 2025, Manuscript No. DMCR-24-33332 (R); **Published:** August 10, 2025, DOI: 10.35248/2684-124X.25.10.4.007

Abstract

Background: Onychomycosis (OM) is a chronic nail disorder frequently encountered by healthcare providers. Yeasts account for only 2-10% of fungal nail infections. Predisposing factors for OM include trauma, diabetes, immunosuppression, nail psoriasis, old age, exposure to humid environments, occupations involving frequent travel, and handwashing.

Objective: To report a rare case of *Candida albicans* nail infection in an immunocompetent 22-year-old Saudi female, a condition that is not commonly seen.

Case Presentation: We present a case of a 22-year-old, otherwise healthy, Saudi female who presented to our dermatology clinic with a nail lesion lasting nearly two months. Physical examination revealed single, unilateral total nail dystrophy with subungual hyperkeratosis of the ring finger, associated with a hyperpigmented plaque with fine white scales. Surprisingly, culture isolated *Candida albicans*, leading us to initiate treatment with itraconazole, dosed as 200 mg BID for one week, followed by a three-week break, in a pulse-dosing regimen. The nail showed dramatic improvement with this regimen during each monthly visit.

Conclusion: *Candida* onychomycosis rarely occurs in immunocompetent adults. Proper history and physical examination are essential for making the correct diagnosis. Culture is the definitive method for identifying the causative microorganism. Nail hygiene and antifungal treatments are the mainstays of treatment.

Keywords: Onychomycosis • *Candida albicans* • Nail infection

Introduction

Background: Onychomycosis (OM) is a chronic nail disorder frequently encountered by healthcare providers [1]. This condition is persistent and not self-limiting; its main signs include nail thickening, brittleness, discoloration, and separation of the nail plate. Yeasts account for only 2–10% of fungal nail infections [2]. Predisposing factors for OM include trauma, diabetes, immunosuppression, nail psoriasis, old age, exposure to humid environments, and occupations involving frequent travel and frequent handwashing [3].

Objective: To report a rare case of *C. albicans* nail infection in an immunocompetent 22-year-old Saudi female, a condition that is not commonly seen.

Case Representation

We present a case of a 22-year-old, otherwise healthy, Saudi female who presented to our dermatology clinic with a nail lesion lasting nearly two months. She denied any history of recent nail trauma, manicure, hepatitis, or travel to areas where hepatitis is endemic. She has no pets, does not consume alcohol, and engages in no recreational drug use. No family members reported similar issues, and there was no history of tinea manuum, cruris, or psoriasis. Physical examination revealed single, unilateral total nail dystrophy with subungual hyperkeratosis of the ring finger, associated with a hyperpigmented plaque with fine white scales, with no involvement of other fingers or toenails. The skin, hair, and mucous membranes were clear with no signs of inflammation. Initial Potassium Hydroxide (KOH) examination showed fungal elements but no specific fungal culture results. Assuming a dermatophyte infection, the most common cause of OM, we prescribed terbinafine 250 mg PO daily for three months, with adjuvant topical therapy, but observed no improvement. Subsequently, specific fungal cultures were performed, which surprisingly isolated *Candida albicans*. Consequently, we initiated itraconazole treatment with a 200 mg BID dose for one week, followed by a three-week break, in a pulse-dosing regimen. The nail showed dramatic improvement with each monthly visit. The patient was monitored closely for potential adverse effects, including hepatotoxicity, and her liver function was regularly assessed. She responded well to the treatment, and her nail condition improved significantly (Figure 1).




Figure 1. Candidal onychomycosis showing nail discoloration and periungual inflammation.

Results and Discussion

Although not life-threatening, OM poses a significant public health problem, accounting for more than 50% of all nail infections [4]. Unfortunately, reports on the incidence of OM in Saudi Arabia are sparse. OM can be caused by dermatophytes, non-dermatophytes, and yeasts. Molecular biology has shown that multiple fungal organisms can cause OM [5]. While *Candida* yeasts are relatively harmless, they often present with an underlying systemic disease [6].

A review of the literature shows that various *Candida* species can cause onychomycosis [7]. Diagnosing onychomycosis depends on patient history, clinical examination, direct microscopic investigation, mycological culture, and histopathology. Initially, we suspected onychomycosis caused by dermatophytes, given their commonality based on history and clinical examination. However, it is crucial to recognize that nearly half of nail infections are not mycotic, necessitating mycological testing for accurate diagnosis [8].

Typically, onychomycosis presents with asymptomatic, dry, hyperkeratotic tinea pedis [9]. In addition, *Candida* organisms usually require an immunocompromised host or risk factors such as repeated nail trauma, diabetes, immunosuppression, nail psoriasis, old age, exposure to humid environments, and occupations involving frequent travel and frequent handwashing. In this case, we present *Candida albicans* onychomycosis without risk factors. Most reported cases involve one or two risk factors, while our patient was immunocompetent and reported no risk factors.

Tests typically used to confirm onychomycosis in a clinical setting include Potassium Hydroxide (KOH) preparations, our first step in investigating the patient's lesion. When the patient did not respond to the antifungal treatment, we utilized fungal cultures. Although fungal cultures are specific, they are not very sensitive and were not chosen initially to reduce costs. Additionally, fungal cultures grow slowly, adding time and expenses to the workup.

Restoring immune defenses, eliminating fungi, using appropriate drug therapy, and improving nail hygiene by addressing predisposing factors are critical in managing candidal onychomycosis.

Conclusion

Candida onychomycosis rarely occurs in immunocompetent adults. Proper history and physical examination are essential for making the correct diagnosis. Culture is the definitive method for identifying the causative microorganism. Nail hygiene and antifungal treatments are the mainstays of treatment.

Patient Consent

Written informed consent was obtained from the patient for publication and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal upon request.

Authors' Contribution

All authors contributed to the preparation of this article. The first author performed the final proofreading.

Conflict of Interest

None declared.

Financial Support and Sponsorship

None.

References

1. Gupta, A.K., et al. "Global prevalence of onychomycosis in general and special populations: An updated perspective." *Mycoses* 67.4 (2024): e13725.
2. Zhang, M.J., et al. "Onychomycosis caused by *Pichia guilliermondii*: A case report and mini-review." *Med Mycol Case Rep* 27 (2020): 72-76.
3. Jazdarehee, A., et al. "Transmission of Onychomycosis and Dermatophytosis between Household Members: A Scoping Review." *J Fungi (Basel)* 8.1 (2022): 60.
4. Ge, G., et al. "Different toenail onychomycosis due to *Rhodotorula mucilaginosa* and *Candida parapsilosis* in an immunocompetent young adult." *Med Mycol Case Rep* 24 (2019): 69-71.
5. Lim, S.S., et al. "Diagnosis of Onychomycosis: From Conventional Techniques and Dermoscopy to Artificial Intelligence." *Front Med* 8 (2021): 637216.
6. Bertolini, M., & Dongari-Bagtzoglou, A. "The Relationship of *Candida albicans* with the Oral Bacterial Microbiome in Health and Disease." *Adv Exp Med Biol* 1197 (2019): 69-78.
7. Hashemi, S.E., et al. "Species distribution and susceptibility profiles of *Candida* species isolated from vulvovaginal candidiasis, emergence of *C. lusitanae*." *Curr Med Mycol* 5.4 (2019): 26-34.
8. Medina, N., et al. "Diagnostic Mycology Laboratories Should Have a Central Role for the Management of Fungal Disease." *J Fungi (Basel)* 8.12 (2022): 1285.
9. Daggett, C., et al. "Onychomycosis in Athletes." *Am J Clin Dermatol* 20.5 (2019): 691-698.