

Changes in the GLUT4 Expression by Acute Exercise, Exercise Training and Detraining in Experimental Models

Alexandre M Lehnen^{1,2}, Kátia De Angelis³, Melissa Medeiros Markoski¹ and Beatriz D'Agord Schaan^{1,2*}

¹Laboratório de Experimentação Animal e Laboratório de Cardiologia Molecular e Celular do Instituto de Cardiologia do RS/Fundação Universitária de Cardiologia do Rio Grande do Sul, Porto Alegre, Brasil

²Serviço de Endocrinologia, Hospital de Clínicas de Porto Alegre/Universidade Federal do Rio Grande do Sul, Porto Alegre, Brasil

³Universidade Nove de Julho, São Paulo, Brasil

Abstract

There is a direct correlation between an increase in insulin sensitivity and increased cell surface GLUT4 content. Acute exercise promotes glucose-transport stimulation that is independent of AMPK and CaMKII insulin-signaling. In turn, post-exercise glucose uptake occurs through changes in components of the insulin signaling cascade involving alterations in downstream mediators such as TBC1D1, TBC1D4/AS160 and p38 MAPK. However, the effects of acute exercise can be reversed within 18–24 hours and appear to be dependent on muscle glycogen levels. Exercise training results in adaptations that facilitate insulin-mediated glucose uptake and are regulated by different mechanisms. It leads to changes in gene expression and greater blood flow and signaling and changes in GLUT4 protein exocytosis and endocytosis. But when exercise training is discontinued GLUT4 tend to return to baseline levels. We have demonstrated in our laboratory that one-week detraining is sufficient to reduce GLUT4 in the heart and adipocytes while this same effect was seen in the gastrocnemius muscle within 2 weeks of training. The present study aimed to review how acute exercise, exercise training, and detraining affect mainly GLUT4 translocation to the insulin-sensitive cell surface.

Keywords: Glucose transporter type 4, Acute exercise, Exercise training, Insulin resistance

Introduction

Glucose transporter 4 (GLUT4) is mainly expressed in insulin-sensitive cells such as adipose tissue and skeletal muscle cells and cardiomyocytes [1]. The main function of this protein is to facilitate glucose uptake into these cells and maintain control of blood glucose levels. GLUT4 protein in the basal state is stored in intracellular vesicles and their translocation to the plasma membrane occurs mainly by insulin action [2] or through insulin-independent pathway during muscle contraction. Insulin resistance may result from impaired insulin signal transduction leading to decreased GLUT4 translocation [3,4] and/or diminished capacity for GLUT4 synthesis [5]. In addition, other factors may also contribute to insulin resistance including reduced blood flow and muscle mass, as well as changes in the proportion of muscle fiber types and in intramuscular oxidative pathways [6].

Acute exercise leads to increased glucose transport even in the presence of very low levels of circulating insulin [7]. During muscle contraction two mechanisms contribute to increasing GLUT4 translocation to the cell surface: activation of the 5'-adenosine monophosphate-activated protein kinase (AMPK) [8] and calcium/calmodulin-dependent protein kinase (CaMK) II [9]. When the acute effects of exercise on glucose transport have disappeared, there are changes in insulin sensitivity [10] that appear to be dependent on muscle glycogen levels [11]. In particular, TBC1D1, TBC1D4/AS160 and p38 MAPK that remain phosphorylated for hours after exercise [12,13], probably activated by residual AMPK of muscle contraction, may contribute to increased insulin sensitivity after exercise. However, these effects may be reversed within 18–24 hours [14,15].

Aerobic [16] and/or resistance exercise training [17] are known to improve insulin sensitivity and have other beneficial effects on blood pressure, heart rate, heart rate variability, and chemoreceptor and arterial baroreceptor reflex sensitivity [18,19]. Given these benefits exercise training has been successfully used in the treatment of diabetic patients [20]. This adaptation is a result of marked increase in glucose trans-

port, which is mostly attributed to greater mobilization of GLUT4-containing vesicles to the cell surface through insulin-dependent [21] and insulin-independent pathways [11]. Besides this effect on GLUT4 trafficking, exercise training also increases transcription factors involved in GLUT4 gene expression and, consequently, increased intracellular GLUT4 stores [22]. Exercise training has also anti-inflammatory effects, which can modify insulin-mediated glucose uptake [23]. However, these benefits are gradually lost after cessation of exercise training. The time course of this phenomenon is still controversial and these changes are affected by several factors. The purpose of this study was to review aspects related to GLUT4 modulation and translocation to the cell surface in response to acute exercise, exercise training, and detraining focusing on studies using experimental models.

GLUT4 and Insulin Resistance

Glucose transport in mammalian tissues occurs primarily by facilitated diffusion, a process that uses a carrier protein for transport of a substrate across a membrane into cells. These facilitative glucose transporters (GLUTs) are a family of proteins that were denominated in chronological order of characterization [24] and are expressed in tissue and cells with different regulatory and kinetic properties that reflect their roles in cellular metabolism.

GLUT4 is the most abundant glucose transporter in the body [1]. In baseline status, i.e. unstimulated cells, GLUT4 is stored intracellularly

***Corresponding author:** Beatriz D'Agord Schaan, Serviço de Endocrinologia, Hospital de Clínicas de Porto Alegre, Centro de Pesquisas, 4º andar, Av. Ramiro Barcelos, 2350, Santa Cecília, 90035-903 Porto Alegre – RS Brasil; E-mail: beatrizschaan@gmail.com

Received June 15, 2012; Accepted July 17, 2012; Published July 21, 2012

Citation: Lehnen AM, Angelis KD, Markoski MM, Schaan BDA (2012) Changes in the GLUT4 Expression by Acute Exercise, Exercise Training and Detraining in Experimental Models. J Diabetes Metab S10:002. doi:[10.4172/2155-6156.S10-002](https://doi.org/10.4172/2155-6156.S10-002)

Copyright: © 2012 Lehnen AM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

in specialized compartments called GLUT4 storage vesicles (GSV), which participate in the GLUT4 cycles to and from the plasma membrane, through slow exocytosis from GSV, fast endocytosis from the cell surface [25-28]. The synthesis of GLUT4 occurs through the expression of the gene SLC2A4 (Solute carrier 2a4 gene that codifies GLUT4 protein) [29] and can determine the amount of protein stored in the GSV.

The main action of insulin is to increase glucose uptake in insulin-sensitive tissues, balancing blood and intra-cellular glucose levels, through increased GLUT4 content at the cell surface. The insulin-mediated glucose uptake is carried out by binding this hormone with its membrane receptors and transmitting its signal to the interior of cells. The insulin-mediated GLUT4 translocation to plasma membrane includes the phosphatidylinositol-3-kinase (PI3K) complex. The c-Cbl-associating protein (CAP/c-Cbl) pathway and its regulation of TC10 downstream also seems involved in GLUT4 translocation to plasma membrane [30]. Decreased insulin action on insulin-sensitive tissues may occur by several mechanisms: low concentration and/or kinase activity of proteins related to insulin signaling, leading to insufficient recruitment of GLUT4 to the plasma membrane, despite normal GLUT4 expression [4]; low capacity of GLUT4 synthesis, even if the rate of translocation of vesicles containing this protein is preserved [5]; and, changes in the rate of GLUT4 exocytosis and endocytosis [31], determined by failure of insulin-derived signals. Thus, insulin resistance is characterized by a reduction of the biological effect of this hormone [32]. Studies have shown a marked relation between insulin resistance and cardiovascular risk factors, among them obesity and sedentary lifestyle [33,34].

The consumption of a high fat diet, even for a short period of time, leads to insulin resistance by reducing the insulin signaling pathway [35]. Thus, experimental models of obesity have been used to elucidate the pathophysiological mechanisms that contribute to the genesis of insulin resistance. High fat diet induced-obesity model has shown that rats subjected to a high fat diet for 3 months developed weight gain, especially epididymal fat, hyperglycemia, hyperinsulinemia and

insulin resistance [36]. The decrease of insulin action was related to protein tyrosine phosphatase 1B (PTP1B). Its increased expression and activation inhibits tyrosine phosphorylation of insulin receptors and their substrates. This is reinforced by experiments with PTP1B knockout mice, which showed high tyrosine phosphorylation of insulin receptors and their substrates, and improved insulin sensitivity as compared to wild mice [37].

Decreased insulin-mediated glucose uptake related to obesity is frequently accompanied by an inflammatory state, characteristic of this condition [38]. The increase in adipocyte size is associated with the release of free fatty acids from the adipose tissue and increased production of reactive oxygen species [39], which may promote the production and release of proinflammatory cytokines such as interleukin-6 (IL-6), serum amyloid A protein (SAA) and monocyte chemoattractant protein-1 (MCP-1) [40]. Cytokines released into the bloodstream act as mediators in the migration of monocytes into the adipose tissue [41]. After migration, these cells differentiate into macrophages, which release cytokines, especially tumor necrosis factor- α (TNF- α) [42]. When the signaling pathway of TNF- α is activated, intermediate substrates, like serine kinase c-Jun NH2-terminal (JNK) influence the phosphorylation of IRS1, decreasing the insulin signal transduction [43]. Increased TNF- α expression leads to decreased phosphorylation of insulin receptors and their substrates, contributing to the genesis of insulin resistance related to obesity [44]. In addition to impaired insulin signal transduction, this cytokine also inhibits the transcription factors of the SLC2A4 gene [42], reducing the expression and, consequently, the intracellular protein stores.

Moreover, there is data suggesting that autonomic changes modulate hormonal and immune function, by inducing release of bioactive molecules which are probably involved in the development of cardio metabolic profile changes [45,46]. In fact, Hellstrom examined evidence that development of a diverse group of diseases, such as diabetes, hypertension, and heart disease, is favored by increased sympathetic neural outflow resulting in endothelial dysfunction, dyslipidemia,




Figure 1: Putative metabolic pathways leading to changes in GLUT4 expression in the skeletal muscle following acute exercise and exercise training. Most GLUT4 is stored in intracellular vesicles (GSV = GLUT4 storage vesicles) in the muscle at rest and their translocation to the plasma membrane occurs mainly by insulin action; after glucose transport, GLUT4 is endocytosed from the cell surface. Muscle contraction leads to an increase in glucose uptake by the working muscle evidenced by gain of GLUT4 at cell surface, which does not involve any signals proximal to the insulin receptor substrate (IRS), phosphatidylinositol 3-kinase (PI3K) or Akt. Instead, this gain occurs by increased AMPK activity that stems from increased ATP-turnover. CaMK protein is also involved in GLUT4 trafficking, although it is unclear whether CaMK is AMPK-dependent or not. Together, these mechanisms increase cellular glucose uptake even in the presence of very low circulating levels of insulin. When the acute effects of exercise on glucose transport have disappeared there are changes in insulin sensitivity that appear to be dependent on TBC1D1, TBC1D4/AS160 and p38 MAPK, which remain phosphorylated for hours after exercise, probably activated by residual AMPK of muscle contraction. As for chronic effects of exercise, evidence suggests there are adaptations in the pre-translational and post-translational levels, in particular MEF2 and GEF, which was shown to increase intracellular GLUT4 and improve insulin sensitivity.

inflammation and insulin resistance [47]. In female fructose fed rats we observed increased arterial pressure and a positive correlation between insulin resistance and cardiac vagal effect attenuation [48]. Recently, we tested the time course of the effect of fructose given in the drinking water in mice, showing: 1) insulin resistance, increased plasma levels of cholesterol, triglycerides and leptin after 60 days of fructose; 2) increase in systolic and mean arterial pressure associated with cardiac and vascular sympathetic increased modulation and spontaneous baroreflex attenuation from day 15 of fructose. The key finding was that dysfunction of cardiovascular autonomic control occurred prior to any metabolic changes [49].

In fact, besides the impairment of extra and intracellular glucose homeostasis, insulin resistance is commonly associated with arterial hypertension and is *per se* a predictor of cardiovascular events [50]. Katayama et al. (1997) showed impaired glucose tolerance in spontaneously hypertensive rats (SHR). Interestingly, GLUT4 in the plasma membrane of the gastrocnemius was elevated in SHR, as compared to control rats (Wistar). However, with 20 weeks of age, SHR normalized glucose tolerance and GLUT4 expression, as compared with Wistar rats. It is possible that the impaired glucose tolerance of SHR have other causes not related to GLUT4, or at least, that GLUT4 is not the only factor contributing to metabolic abnormalities seen in this animal model [51]. This may involve the transport of glucose in the plasma membrane and not the increased levels of GLUT4 at the cell surface. This phenomenon may be explained in part by glucose not binding to GLUT4 on the cell surface, slowing its entry into these cells [52].

Effects of Acute Exercise and Training on GLUT4

Exercise can induce acute metabolic benefits that are different from those induced by training (Figure 1). An acute bout of exercise, through muscle contraction, requires several metabolic changes to supply adenosine triphosphate (ATP), including the use of muscle glycogen. The energy demands of an acute exercise require increased glucose uptake in muscle cells, mainly to muscular glycogen resynthesizing. Thus, acute exercise leads to changes in glucose uptake that occur through insulin-independent pathways [53,54] to ensure an adequate glucose supply to muscle cells. Moreover, acute exercise induces adjustments in the GLUT4 cycling as increased GLUT4 exocytosis and/or lower GLUT4 endocytosis [55,56].

It was observed following the acute effects of exercise a persistent increase in glucose transport for hours that appears to be dependent on the working muscle, [57]. This effect is associated to increased GLUT4 translocation, probably stimulated by downstream TBC1D1 and TBC1D4/AS160 phosphorylation, which is activated by the remaining AMPK activity of muscle contraction [12,13]. Increased glucose uptake after muscle contraction can also be explained by an acute increase in p38 MAPK activation, which remained increased even 3 h after muscle contraction in the soleus and epitrochlearis muscles of rats [58]. Together, TBC1D1, TBC1D4/AS160 and p38 MAPK remain phosphorylated for hours after exercise and may contribute to increased insulin sensitivity after exercise.

In contrast, adaptations in the pre-translational and post-translational levels are observed as chronic effects of exercise. Hence, over time, it leads to enhanced insulin action, either increased expression and activity of insulin-signaling protein kinases and other insulin-independent pathways such as increased GLUT4 transcription factors, resulting in an increase in intracellular GLUT4 stores.

Acute Effects of Exercise

Some studies suggest there are different intracellular pools of GLUT4, one stimulated by insulin and one stimulated by exercise [59,60]. Furthermore, studies in insulin receptor knockout mice showed exercise-mediated increase in glucose uptake and glycogen synthase activity *in vivo* as compared to animals in the control group [61]. These facts supports the idea that the acute exercise can activate molecular pathways of GSV mobilization that are not, at least in part, insulin-dependent. In insulin-independent pathways, two proteins have a major role in the mobilization of GLUT4: AMPK and CaMKII.

AMPK is activated by AMP-binding resulting from an increase in ATP-turnover and evidence from experiments using an AMP-mimetic compound TBC1D1, 5-aminimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR), shows that AMPK may be sufficient to increase glucose transport [62,63], and this effect is related to muscle fiber type [64,65]. Kurth et al. (1999) have showed increased glucose uptake observed with AMPK activation by AICAR in perfused rat hindlimb muscles is due to an increase in the translocation of GLUT4 to surface membranes [8].

One of the mechanisms for GLUT4 trafficking may be the interaction of AMPK with AS160, a protein of the PI3K pathway [66]. The administration of AICAR was found to be associated with increased AS160 phosphorylation together with increased AMPK activity. In contrast, after the administration of Wortmannin, a PI3K inhibitor, AS160 activity was completely inhibited [67]. The same was evidenced in other studies [68]. Thus, it is admissible to think that acute exercise-induced increased AMPK activity can be involved with AS160 phosphorylation, and in this case, it is independent of the insulin signaling.

Besides the acute effects of exercise on AMPK-induced GLUT4 trafficking, a positive correlation was observed between acute exercise-induced increase in CaMKII levels and glucose transport in muscle cells [69]. Evidence obtained using subcontraction concentrations of caffeine, which release Ca^{2+} from the sarcoplasmic reticulum and activate CaMKII, has shown that inhibition of this protein prevents an increase in Ca^{2+} -induced glucose transport [70,71]. Although studies [72,73] have shown that caffeine-stimulated glucose transport is highly AMPK-dependent, there is evidence demonstrating that contraction-induced skeletal muscle glucose uptake involving Ca^{2+} /calmodulin-dependent protein kinase is independent of AMPK [74]. Witczak et al. later demonstrated using a CaMKII inhibitory peptide transfected into tibialis anterior muscles by *in vivo* electroporation that this peptide did not either change GLUT4 expression or impair contraction-induced increases in the phosphorylation of AMPK or TBC1D1 and TBC1D4 on AS160 phosphorylation [75]. This evidence supports that CaMKII plays a critical role in the regulation of contraction-induced glucose uptake.

Other proteins such as aPKC (atypical PKC) have been investigated as modulators of GLUT4 translocation in response to acute exercise [13]. Chen et al. (2002) found that glucose uptake in L6 muscle cells (cultured muscle fibers) resulted from protein kinase activation including the isoforms PKC- ζ and PKC- λ among others. A possible explanation for this association would be an interaction of PKC with the AMPK signaling pathway that in turn stimulates GLUT4 translocation [76].

Nitric oxide (NO) has also been investigated as a potential factor to induce increased glucose uptake into muscle cells as an adjustment to acute exercise. Roberts et al. (1997) used nitro-L-arginine methyl ester (L-NAME), an inhibitor of NO, in the gastrocnemius muscle of

Sprague-Dawley rats undergo acute exercise and acute exercise plus L-NAME. After the one bout of exercise (treadmill exercise for 45 min at high intensity), a group received L-NAME in the drinking water (1g/mL). The group that only exercised without L-NAME showed increased glucose uptake and GLUT4 in the plasma membrane of the gastrocnemius muscle as compared to the control group. The group that exercised plus L-NAME did not show any changes in either glucose uptake or the amount of GLUT4 at the cell surface compared to controls [77]. One possible explanation for the effect of NO on insulin-mediated glucose uptake would be an interaction with the AMPK pathway [78], in particular $\alpha 2$ -AMPK site that seems to have a more relevant role in the regulation of glucose transport than $\alpha 1$ -AMPK [79,80].

Another aspect may be related to S-nitrosation of the proteins involved in the insulin signaling cascade, which reduces insulin action [81-82]. This hypothesis is supported by experiments in Wistar rats submitted to high-fat diet-induced obesity, showing high levels of iNOS and S-nitrosation of IR β subunit, IRS1 and Akt, changes that were reversed by acute exercise. The improvements were ascribed to increased AMPK activity that negatively modulates iNOS levels and subsequent S-nitrosation of the proteins involved in insulin signaling [83].

Thus, it seems clear that several insulin-independent pathways are activated in response to acute exercise. The major pathways are those mediated by AMPK protein activation and are dependent on the muscle fiber type.

Chronic Effects of Exercise

Exercise-induced chronic effects on GLUT4 occur primarily through pre-translational mechanisms, which favor the increase of intracellular stores of GLUT4 protein. Besides this beneficial adaptation, exercise training also determines changes in molecular pathways that induce GLUT4 translocation. The changes of exercise training can also be observed in the total GLUT4 protein levels, i.e. GLUT4 expression at cell surface enriched of microsomal fraction.

In this context, Neuffer et al. (1992) studied Wistar rats that were submitted to exercise training on a treadmill for 1 day, 1 week or 6 weeks (1.9 km/h, 2 hours a day, 6 days a week). Biopsy samples of soleus and vastus lateralis muscles (red and white fibers) were examined and compared with those of animals that did not exercise. There was no difference in the amount of GLUT4 in the muscles after 1 day or 1 week of exercise, suggesting that exercise load was insufficient. Exercise training for 6 weeks resulted in increased levels of GLUT4 in the plasma membrane of soleus (1.4 times) and in the oxidative muscle, but not in the glycolytic fibers of the vastus lateralis muscle (1.7 times). Probably cell requirements related to cellular oxidative capacity modulates GLUT4 translocation, as mentioned above. Another explanation would be that dynamic aerobic training was not enough to exercise or even activate glycolytic fibers of the vastus lateralis muscle [84].

To study different types of exercise training, 6-week-old Sprague-Dawley rats were submitted to resistance training (3 sets of lifting of 75% one-repetition maximum, 10 repetitions, 3 days a week for 12 weeks) or aerobic training (1.9 km/h, 15% incline, 45 min a day, 3 days a week for 12 weeks) and compared with a sedentary group. After training periods, rats received an insulin infusion and samples of the soleus, plantar and oxidative and glycolytic fibers of the gastrocnemius and quadriceps muscles were collected. There was a similar increase in glucose uptake in the groups submitted to resistance and aerobic training and both had higher glucose uptake than the sedentary group.

There was a higher rate of glucose transport in rats subjected to the resistance training, probably due to an increase in the amount of GLUT4 at the cell surface [17].

Our group found a similar result in SHR and WKY rats. There was no difference in GLUT4 content in the plasma membrane of the heart, gastrocnemius muscle and epididymal fat of SHR submitted to exercise training (treadmill, 1h/day, 5 days/week for 10 weeks) as compared to WKY rats [85]. Song et al. (1998) reported no difference in GLUT4 expression in samples of gastrocnemius muscle in response to swim training for 4 weeks between stroke-prone spontaneously hypertensive (SHRSP) rats, which are characteristically resistant to insulin, and WKY rats [86].

Luciano et al. (2002) described improved glucose tolerance and increased total GLUT4 expression in the gastrocnemius muscle of Wistar rats submitted to a 6 week regimen of swimming [21]. These changes were attributed to an increase in insulin signaling due to greater phosphorylation of IRS1 and IRS2 substrates. PI3K kinase activity associated with IRS1 and IRS2 as well as serine phosphorylation of Akt protein also increased in trained animals, as compared to sedentary ones. Chibalin et al. (2000) reported that Wistar rats submitted to swim training had increased GLUT4 expression in the epitrochlearis muscles. These adaptations resulted in part from increased tyrosine phosphorylation of insulin receptors and its IRS1 and IRS2 substrates, as well as its association with PI3K protein [87]. Akt activity also increased after exercise training. On the other hand, only the expression levels of insulin receptors were high. Interestingly, the expression of IRS1 substrate after 5 days of swim training was reduced, despite an increased IRS1 phosphorylation; as IRS1 and IRS2 responded differently to exercise training; these two molecules may have different roles and regulations in insulin signaling.

Data from our group showed that chronic treatment with L-NAME induced not only hemodynamic impairment but also insulin resistance, which was not reversed after exercise training at the baseline [88]. These results suggest an important role of NO not only in the development of insulin resistance at baseline, but also in adaptive responses to exercise training.

Another hypothesis for increased GLUT4 at the cell surface in response to exercise training may be effects of galanin. Galanin is a neuroendocrine peptide and an important hormone in insulin sensitivity modulation [89]. He et al. (2011) used an antagonist of galanin (M35) and a regimen of 4-week swim training (60 min per day) to determine whether increased galanin would elevate GLUT4 concentration in the plasma membrane of hind-limb skeletal-muscle of streptozotocin-induced diabetic rats. It was shown that plasma galanin levels after swim training were higher as compared with the sedentary control group. The antagonization of galanin reduced glucose disappearance rate of euglycemic hyperinsulinemic clamp tests when compared with diabetic controls, but swimming enhanced insulin sensitivity in all trained groups. Moreover, M35 treatment reduced GLUT4 concentration and mRNA levels compared with the diabetic control group. In contrast, all trained groups showed an increase of the GLUT4 mRNA expression and GLUT4 protein level of hind-limb skeletal-muscle [90]. Considering these results, endogenous galanin may enhance glucose disappearance rate by increased GLUT4 content in the skeletal muscle's plasma membrane.

Finally, the increase in GLUT4 at the cell surface in response to exercise training may also suggest an effect on GLUT4 mRNA. SLC2A4 gene transcription is activated by two main factors: myocyte enhancer

factor 2 (MEF2) [91] and GLUT4 enhancer factor (GEF) [92]. The rest of MEF2 is related to class II histone deacetylase 5 gene isoform (HDAC5), a molecule that represses SLC2A4 gene transcription. On the other hand, AMPK protein activation through muscle contraction requires HDAC5 phosphorylation resulting in MEF2 release [93]. Hence, another interesting aspect of increased AMPK activity in response to exercise, although not directly related to GLUT4 trafficking to the plasma membrane, is its interaction with transcriptional activation of SLC2A4 gene via MEF2D activity [22], which could increase intracellular GLUT4 stores. However, it is unclear whether a single bout of exercise is strong enough to functionally affect GLUT4 intracellular stores or it is a chronic positive effect of exercise training.

Collectively, data suggest that dynamic aerobic training and/or resistance lead to increased GLUT4 expression levels in the insulin-sensitive cells. This fact increases the uptake of insulin-mediated glucose, providing an appropriate glycemic balance between tissue and plasma in the resting state.

Effects of Exercise Detraining on GLUT4

Although exercise training is beneficial, increasing insulin sensitivity, this adaptation is transient. When this practice is stopped or the stimulus of exercise training is not sufficient to lead to further physiological adaptations, the trend is for the transport of glucose to return back to baseline values [94,95]. It has been demonstrated that the metabolic improvements decline with different degrees during detraining time course [96]. However, most of the studies were made with athletes or healthy subjects and there are only few works focusing on the association of detraining and disease.

Furthermore, in our laboratory, using SHR, we found that after 1 and 2 weeks of exercise detraining, animals remain with the improvement in insulin sensitivity (whole-body insulin sensitivity measured by the insulin tolerance test – ITT) and lower blood pressure levels determined by exercise training on the treadmill for 10 weeks. However, the reversal of the increased expression of GLUT4 occurred after one week of detraining in the heart and adipose tissue, and after two weeks in skeletal muscle (gastrocnemius) [95]. Other authors showed that one week is sufficient to reverse the benefits on GLUT4 in skeletal muscle [84]. However, the difference might be the type of fiber used. This hypothesis suggests that different molecular mechanisms governing the process of exercise detraining is mediated by a tissue-specific modulation of expression of GLUT4. Neufer et al. (1992) analyzed samples of the soleus and vastus lateralis (red fibers) which are essentially oxidative [97], while our research was performed with samples of the gastrocnemius with no separation of white and red fibers. Other explanation would be that the effect of exercise detraining is dependent on the extension of the exercise training. Our training regimen was based on 5 days per week for 10 weeks (50 sessions). Neufer et al. (1992) used 6 days a week for 6 weeks (36 sessions).

To study the effects of exercise detraining on insulin receptors the epitrochlearis muscle of Fischer rats was analyzed 29 and 53 hours after cessation of 3-weeks of voluntary wheel running. GLUT4 protein levels in the plasma membrane of epitrochlearis muscle returned to sedentary levels (reduction of 29%) 53 hours after the cessation of the physical activity. This fact could be partially explained by decrease in tyrosine phosphorylation and protein level of IR (β -subunit) as well as Akt phosphorylation activity. All these variables also returned to sedentary levels after the same period of detraining (53 hours) [98].

Reynolds et al. (2000) observed the effects of swimming (5 days or

5 weeks) or treadmill (5 weeks), followed by 1 or 2 days of detraining period [99]. The amount of GLUT4 in the plasma membrane remained high during 24 hours after training in all groups trained, but returned to baseline levels 48 hours after the last bout of exercise, regardless of the period or exercise regimen. One possible explanation is an adaptation of the GLUT4 half-life, which can be regulated by pre- and post-translational mechanisms. The increased half-life also results from a decrease in the rate of degradation of the protein; in this case, the rate of reversal of the adaptation would occur more rapidly than its development [94].

Mostarda et al. (2009) evaluated the effect of 3-week detraining after 10 weeks of training in streptozotocin-induced diabetic rats. Despite evidence showing that GLUT4 expression returns to pre-training levels within 48 hours to 1 week [84,95,99], they found that, after 3 weeks of detraining, glucose levels remained similar to those in the trained group [100]. These data show that glucose clearance cannot be explained only by GLUT4 expression levels as it was demonstrated.

Although the biological effects of exercise detraining on glucose metabolism are evident, the molecular mechanisms of this down-regulation have not been totally understood, especially on insulin-independent pathways. Studies are needed to determine the involvement and time-course of these mechanisms to lead to effective prevention management and treatment of insulin resistance and its consequences.

Conclusions

Based on the assumptions presented, we conclude that: 1) acute exercise increases GLUT4 expression by parallel insulin signaling pathways; 2) most mechanisms implicated involve activation of AMPK and/or CaMKII; 3) AMPK-mediated glucose uptake is higher in fast-twitch muscle fibers; 4) increased glucose uptake following acute exercise may also involve changes in GLUT4 exocytosis and endocytosis rates; 5) improved insulin sensitivity following acute exercise results from adaptations linked to TBC1D1, TBC1D4/AS160 and p38 MAPK, which remain active for hours after muscle contraction cessation; and finally 6) adaptations in the pre-translational and post-translational levels are regarded as chronic effects of exercise. On the other hand, cessation of exercise training leads to a decrease in the amount of GLUT4 in the plasma membrane in animal models. We conclude that: 1) training-induced beneficial effects on GLUT4 expression can be reversed within 48 h to 1 week following training cessation; 2) the effects of detraining on GLUT4 expression may involve pre- and post-translational mechanisms; and 3) glucose clearance cannot be explained only by GLUT4 levels, at least not in insulin-resistant rats.

This review shows the importance of acute and chronic exercise in changes in GLUT4 expression by insulin-independent pathways, which remain intact even in individuals with insulin resistance.

Competing Interests

The authors declare that they have no competing interests.

References

1. Ploug T, Van Deurs B, Ai H, Cushman SW, Ralston E (1998) Analysis of GLUT4 distribution in whole skeletal muscle fibers: identification of distinct storage compartments that are recruited by insulin and muscle contractions. *J Cell Biol* 142: 1429-1446.
2. Carvalho E, Jansson PA, Nagaev I, Wenthzel AM, Smith U (2001) Insulin resistance with low cellular IRS-1 expression is also associated with low GLUT4 expression and impaired insulin-stimulated glucose transport. *FASEB J* 15: 1101-1113.
3. Garvey W, Malanu L, Zhu JH, Brechtel-Hook G, Wallace P, et al. (1998)

- Evidence for defects in the trafficking and translocation of GLUT4 glucose transporters in skeletal muscle as a cause of human insulin resistance. *J Clin Invest* 101: 2377-2386.
4. Bjornholm M, Zierath JR (2005) Insulin signal transduction in human skeletal muscle: identifying the defects in Type II diabetes. *Biochem Soc Trans* 33: 354-357.
 5. Machado UF, Shimizu I, Saito M (1994) Reduced content and preserved translocation of glucose transporter (GLUT 4) in white adipose tissue of obese mice. *Physiol Behav* 55: 621-625.
 6. LaMonte MJ, Blair SN, Church TS (2005) Physical activity and diabetes prevention. *J Appl Physiol* 99:1205-1213.
 7. Wasserman DH, Mohr T, Kelly P, Lacy DB, Bracy D (1992) Impact of insulin deficiency on glucose fluxes and muscle glucose metabolism during exercise. *Diabetes* 41: 1229-1238.
 8. Kurth-Kraczek EJ, Hirshman MF, Goodyear LJ, Winder WW (1999) 5'AMP-activated protein kinase activation causes GLUT4 translocation in skeletal muscle. *Diabetes* 48: 1667-1671.
 9. Smith JA, Collins M, Grobler LA, Magee CJ, Ojuka EO (2007) Exercise and CaMK activation both increase the binding of MEF2A to the Glut4 promoter in skeletal muscle in vivo. *Am J Physiol Endocrinol Metab* 292: 413-420.
 10. Richter EA, Garetto LP, Goodman MN, Ruderman NB (1982) Muscle glucose metabolism following exercise in the rat: increased sensitivity to insulin. *J Clin Invest* 69: 785-793.
 11. Ren JM, Semenkovich CF, Gulve EA, Gao J, Holloszy JO (1994) Exercise induces rapid increases in GLUT4 expression, glucose transport capacity, and insulin-stimulated glycogen storage in muscle. *J Biol Chem* 269: 14396-143401.
 12. Sakamoto K, Holman GD (2008) Emerging role for AS160/TBC1D4 and TBC1D1 in the regulation of GLUT4 traffic. *Am J Physiol Endocrinol Metab* 295: E29-E37.
 13. Maarbjerg SJ, Sylow L, Richter EA (2011) Current understanding of increased insulin sensitivity after exercise - emerging candidates. *Acta Physiol (Oxf)* 202: 323-335.
 14. Kuo CH, Hwang H, Lee MC, Castle AL, Ivy JL (2004) Role of insulin on exercise-induced GLUT-4 protein expression and glycogen supercompensation in rat skeletal muscle. *J Appl Physiol* 96: 621-627.
 15. Host HH, Hansen PA, Nolte LA, Chen MM, Holloszy JO (1998) Glycogen supercompensation masks the effect of a training-induced increase in GLUT4 on muscle glucose transport. *J Appl Physiol* 85: 133-138.
 16. Rodnick KJ, Holloszy JO, Mondon CE, James DE (1990) Effects of exercise-training on insulin-regulatable glucose-transporter protein levels in rat skeletal muscle. *Diabetes* 39: 1425-1429.
 17. Yaspelkis BB 3rd, Singh MK, Trevino B, Krisan AD, Collins DE (2002) Resistance training increases glucose uptake and transport in rat skeletal muscle. *Acta Physiol Scand* 175: 315-323.
 18. Harthmann AD, De Angelis K, Costa LP, Senador D, Schaan BD, et al. (2007) Exercise training improves arterial baro- and chemoreflex in control and diabetic rats. *Auton Neurosci* 133: 115-120.
 19. Loimaala A, Huikuri HV, Koobi T, Rinne M, Nenonen A, et al. (2003). Exercise training improves baroreflex sensitivity in type 2 diabetes. *Diabetes* 52: 1837-1842.
 20. Umpierre D, Ribeiro PA, Kramer CK, Leitao CB, Zucatti AT, et al. (2011) Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 305: 1790-1799.
 21. Luciano E, Carneiro EM, Carvalho CR, Carvalheira JB, Peres SB, et al. (2002) Endurance training improves responsiveness to insulin and modulates insulin signal transduction through the phosphatidylinositol 3-kinase/Akt-1 pathway. *Eur J Endocrinol* 147: 149-157.
 22. Lima GA, Anhê GF, Giannocco G, Nunes MT, Correa-Giannella ML, et al. (2009) Contractile activity per se induces transcriptional activation of SLC2A4 gene in soleus muscle: involvement of MEF2D, HIF-1a, and TRalpha transcriptional factors. *Am J Physiol Endocrinol Metab* 296: E132-E138.
 23. Geffken DF, Cushman M, Burke GL, Polak JF, Sakkinen PA, et al. (2001). Association between physical activity and markers of inflammation in a healthy elderly population. *Am J Epidemiol* 153: 242-250.
 24. Uldry M, Thorens B (2004) The SLC2 family of facilitated hexose and polyol transporters. *Pflugers Arch* 447: 480-489.
 25. Satoh S, Nishimura H, Clark AE, Kozka IJ, Vannucci SJ, et al. (1993) Use of bismannose photolabel to elucidate insulin-regulated GLUT4 subcellular trafficking kinetics in rat adipose cells. Evidence that exocytosis is a critical site of hormone action. *J Biol Chem* 268: 17820-17829.
 26. Li D, Randhawa VK, Patel N, Hayashi M, Klip A (2001) Hyperosmolarity reduces GLUT4 endocytosis and increases its exocytosis from a VAMP2-independent pool in I6 muscle cells. *J Biol Chem* 276: 22883-22891.
 27. Klip A (2009) The many ways to regulate glucose transporter 4. *Appl Physiol Nutr Metab* 34: 481-487.
 28. Klip A, Schertzer JD, Bilan PJ, Thong F, Antonescu C (2009) Regulation of glucose transporter 4 traffic by energy deprivation from mitochondrial compromise. *Acta Physiol (Oxf)* 196: 27-35.
 29. Joost HG, Bell GI, Best JD, Birnbaum MJ, Charron MJ, et al. (2002) Nomenclature of the GLUT/SLC2A family of sugar/polyol transport facilitators. *Am J Physiol Endocrinol Metab* 282: E974-E976.
 30. Chiang SH, Baumann CA, Kanzaki M, Thurmond DC, Watson RT, et al. (2001) Insulin-stimulated GLUT4 translocation requires the CAP-dependent activation of TC10. *Nature* 410: 944-948.
 31. Foster LJ, Li D, Randhawa VK, Klip A (2001) Insulin accelerates inter-endosomal GLUT4 traffic via phosphatidylinositol 3-kinase and protein kinase B. *J Biol Chem* 276: 44212-44221.
 32. Shaw DI, Hall WL, Williams C (2005) Metabolic Syndrome: What is it and What are the Implications? *Proc Nutr Soc* 64: 349-357.
 33. Bergman RN, Kim SP, Hsu IR, Catalano KJ, Chiu JD, et al. (2007) Abdominal obesity: role in the pathophysiology of metabolic disease and cardiovascular risk. *Am J Med* 120: S3-S8.
 34. DeFronzo RA, Ferrannini E (1991) Insulin resistance. A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diabetes Care* 14: 173-194.
 35. Clegg DJ, Gotoh K, Kemp C, Wortman MD, Benoit SC, et al. (2011) Consumption of a high-fat diet induces central insulin resistance independent of adiposity. *Physiol Behav* 103: 10-16.
 36. Ropelle ER, Pauli JR, Prada PO, de Souza CT, Picardi PK, et al. (2006) Reversal of diet-induced insulin resistance with a single bout of exercise in the rat: the role of PTP1B and IRS-1 serine phosphorylation. *J Physiol* 577: 997-1007.
 37. Elchebly M, Payette P, Michaliszyn E, Cromlish W, Collins S, et al. (1999) Increased insulin sensitivity and obesity resistance in mice lacking the protein tyrosine phosphatase-1B gene. *Science* 283: 1544-1548.
 38. Nguyen MT, Satoh H, Favelyukis S, Babendure JL, Imamura T et al. (2005) JNK and tumor necrosis factor- α mediate free fatty acid-induced insulin resistance in 3T3-L1 adipocytes. *J Biol Chem* 280: 35361-35371.
 39. Tchoukalova YD, Sarr MG, Jensen MD (2004) Measuring committed preadipocytes in human adipose tissue from severely obese patients by using adipocyte fatty acid binding protein. *Am J Physiol Regul Integr Comp Physiol* 287: 1132-1140.
 40. Takahashi K, Mizuarai S, Araki H, Mashiko S, Ishihara A, et al. (2003) Adiposity elevates plasma MCP-1 levels leading to the increased CD11b-positive monocytes in mice. *J Biol Chem* 278: 46654-46660.
 41. Curat CA, Miranville A, Sengenès C, Diehl M, Tonus C, et al. (2004) From blood monocytes to adipose tissue-resident macrophages: induction of diapedesis by human mature adipocytes. *Diabetes* 53: 1285-1292.
 42. Stephens JM, Pekala PH (1991) Transcriptional repression of the GLUT4 and C/EBP genes in 3T3-L1 adipocytes by tumor necrosis factor- α . *J Biol Chem* 266: 21839-21845.
 43. Hirosumi J, Tuncman G, Chang L, Gorgun CZ, Uysal KT, et al. (2002) A central role for JNK in obesity and insulin resistance. *Nature* 420: 333-336.
 44. Hotamisligil GS, Peraldi P, Budavari A, Ellis R, White MF, et al. (1996)

- Spiegelman BM. IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF- α - and obesity-induced insulin resistance. *Science* 271: 665-668.
45. Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, et al. (2000) Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405: 458-462.
46. Van Gaal LF, Mertens IL, De Block CE (2006) Mechanisms linking obesity with cardiovascular disease. *Nature* 444: 875-880.
47. Hellstrom HR (2007) The altered homeostatic theory: A hypothesis proposed to be useful in understanding and preventing ischemic heart disease, hypertension, and diabetes—including reducing the risk of age and atherosclerosis. *Med Hypotheses* 68: 415-433.
48. Brito JO, Ponciano K, Figueroa D, Bernardes N, Sanches IC, et al. (2008) Parasympathetic dysfunction is associated with insulin resistance in fructose-fed female rats. *Braz J Med Biol Res* 41: 804-808.
49. De Angelis K, Senador DD, Mostarda CT, Irigoyen MC, Morris M (2012) Sympathetic Overactivity Precedes Metabolic Dysfunction in a Fructose Model of Glucose Intolerance in Mice. *Am J Physiol Regul Integr Comp Physiol* 302: R950-R957.
50. Sarwar N, Danesh J, Eiriksdottir G, Sigurdsson G, Wareham N, et al. (2007) Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation* 115: 450-458.
51. Katayama S, Inaba M, Maruno Y, Morita T, Awata T, et al. (1997) Glucose intolerance in spontaneously hypertensive and wistar-kyoto rats: enhanced gene expression and synthesis of skeletal muscle glucose transporter 4. *Hypertens Res* 20: 279-286.
52. Collison M, James DJ, Grahan D, Homan GD, Connell JM, et al. (2005) Reduced insulin-stimulated GLUT4 bioavailability in stroke-prone spontaneously hypertensive rats. *Diabetologia* 48: 539-546.
53. Wright DC, Hucker KA, Holloszy JO, Han DH (2004) Ca^{2+} and AMPK both mediate stimulation of glucose transport by muscle contractions. *Diabetes* 53: 330-335.
54. Lira VA, Soltow QA, Long JH, Betters JL, Sellman JE, et al. (2007) Nitric oxide increases GLUT4 expression and regulates AMPK signaling in skeletal muscle. *Am J Physiol Endocrinol Metab* 293: E1062-E1068.
55. Antonescu CN, Diaz M, Femia G, Planas JV, Klip A (2008) Clathrin-dependent and independent endocytosis of glucose transporter 4 (GLUT4) in myoblasts: regulation by mitochondrial uncoupling. *Traffic* 9: 1173-1190.
56. Wijesekara N, Tung A, Thong F, Klip A (2006) Muscle cell depolarization induces a gain in surface GLUT4 via reduced endocytosis independently of AMPK. *Am J Physiol Endocrinol Metab* 290: 1276-1286.
57. Jensen TE, Richter EA (2012) Regulation of glucose and glycogen metabolism during and after exercise. *J Physiol* 590: 1069-1076.
58. Geiger PC, Wright DC, Han DH, Holloszy JO (2005) Activation of p38 MAP kinase enhances sensitivity of muscle glucose transport to insulin. *Am J Physiol Endocrinol Metab* 288: E782-E788.
59. Douen AG, Ramlal T, Rastogi S, Bilan PJ, Cartee GD, et al. (1990) Exercise induces recruitment of the "insulin-responsive glucose transporter". Evidence for distinct intracellular insulin- and exercise-recruitable transporter pools in skeletal muscle. *J Biol Chem* 265: 13427-13430.
60. Coderre L, Kandror KV, Vallega G, Pilch PF (1995) Identification and characterization of an exercise-sensitive pool of glucose transporters in skeletal muscle. *J Biol Chem* 270: 27584-27588.
61. Wojtaszewski JF, Higaki Y, Hirshman MF, Michael MD, Dufresne SD, et al. (1999) Exercise modulates postreceptor insulin signaling and glucose transport in muscle-specific insulin receptor knockout mice. *J Clin Invest* 104: 1257-1264.
62. Jessen N, An D, Lihn AS, Nygren J, Hirshman MF, et al. (2011) Exercise increases TBC1D1 phosphorylation in human skeletal muscle. *Am J Physiol Endocrinol Metab* 301: E164-E171.
63. Witczak CA, Sharoff CG, Goodyear LJ (2008) AMP-activated protein kinase in skeletal muscle: from structure and localization to its role as a master regulator of cellular metabolism. *Cell Mol Life Sci* 65: 3737-3755.
64. Jorgensen SB, Viollet B, Andreelli F, Frosig C, Birk JB, et al. (2004) Knockout of the $\alpha 2$ but not $\alpha 1$ 5'-AMP-activated protein kinase isoform abolishes 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranosidebut not contraction-induced glucose uptake in skeletal muscle. *J Biol Chem* 279: 1070-1079.
65. Wright DC, Geiger PC, Holloszy JO, Han DH (2005) Contraction- and hypoxia-stimulated glucose transport is mediated by a Ca^{2+} dependent mechanism in slow-twitch rat soleus muscle. *Am J Physiol Endocrinol Metab* 288: E1062-E1066.
66. Arias EB, Kim J, Funai K, Cartee GD (2007) Prior exercise increases phosphorylation of Akt substrate of 160 kDa (AS160) in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 292: 1191-1200.
67. Kramer HF, Witczak CA, Fujii N, Jessen N, Taylor EB, et al. (2006) Distinct signals regulate AS160 phosphorylation in response to insulin, AICAR, and contraction in mouse skeletal muscle. *Diabetes* 55: 2067-2076.
68. Treebak JT, Glund S, Deshmukh A, Klein DK, Long YC, et al. (2006) AMPK-mediated AS160 phosphorylation in skeletal muscle is dependent on AMPK catalytic and regulatory subunits. *Diabetes* 55: 2051-2058.
69. Terada S, Muraoka I, Tabata I (2003) Changes in $[\text{Ca}^{2+}]_i$ induced by several glucose transport-enhancing stimuli in rat epitrochlearis muscle. *J Appl Physiol* 94: 1813-1820.
70. Jensen TE, Rose AJ, Hellsten Y, Wojtaszewski JF, Richter EA (2007) Caffeine-induced Ca^{2+} release increases AMPK-dependent glucose uptake in rodent soleus muscle. *Am J Physiol Endocrinol Metab* 293: E286-E292.
71. Wright DC, Hucker KA, Holloszy JO, Han DH (2004) Ca^{2+} and AMPK both mediate stimulation of glucose transport by muscle contractions. *Diabetes* 53: 330-335.
72. Jensen TE, Rose AJ, Hellsten Y, Wojtaszewski JF, Richter EA (2007) Caffeine-induced Ca^{2+} release increases AMPK-dependent glucose uptake in rodent soleus muscle. *Am J Physiol Endocrinol Metab* 293: E286-E292.
73. Abbott MJ, Bogachus LD, Turcotte LP (2011) AMPK $\alpha 2$ deficiency uncovers time dependency in the regulation of contraction-induced palmitate and glucose uptake in mouse muscle. *J Appl Physiol* 111: 125-134.
74. Witczak CA, Fujii N, Hirshman MF, Goodyear LJ (2007) Ca^{2+} /calmodulin-dependent protein kinase kinase- α regulates skeletal muscle glucose uptake independent of AMP-activated protein kinase and Akt activation. *Diabetes* 56: 1403-1409.
75. Witczak CA, Jessen N, Warro DM, Toyoda T, Fujii N, et al. (2010) CaMKII regulates contraction- but not insulin-induced glucose uptake in mouse skeletal muscle. *Am J Physiol Endocrinol Metab* 298: E1150-E1160.
76. Chen HC, Bandyopadhyay G, Sajan MP, Kanoh Y, Standaert M, et al. (2002) Activation of the ERK pathway and atypical protein kinase C isoforms in exercise- and aminoimidazole-4-carboxamide-1- β -D-ribose (AICAR)-stimulated glucose transport. *J Biol Chem* 277: 23554-23562.
77. Roberts CK, Barnard RJ, Scheck SH, Balon TW (1997) Exercise-stimulated glucose transport in skeletal muscle is nitric oxide dependent. *Am J Physiol* 273: E220-E225.
78. Shearer J, Fueger PT, Vorndick B, Bracy DP, Rottman JN, et al. (2004) AMP kinase-induced skeletal muscle glucose but not long-chain fatty acid uptake is dependent on nitric oxide. *Diabetes* 53: 1429-1435.
79. Jessen N, Pold R, Buhl ES, Jensen LS, Schmitz O, et al. (2003) Effects of AICAR and exercise on insulin-stimulated glucose uptake, signaling, and GLUT-4 content in rat muscles. *J Appl Physiol* 94: 1373-1379.
80. Ai H, Ihlemann J, Hellsten Y, Lauritzen HP, Hardie DG, et al. (2002) Effect of fiber type and nutritional state on AICAR- and contraction-stimulated glucose transport in rat muscle. *Am J Physiol Endocrinol Metab* 282: E1291-E1300.
81. Sugita H, Fujimoto M, Yasukawa T, Shimizu N, Sugita M, et al. (2005) Inducible nitric-oxide synthase and NO donor induce insulin receptor substrate-1 degradation in skeletal muscle cells. *J Biol Chem* 280: 14203-14211.
82. Yasukawa T, Tokunaga E, Ota H, Sugita H, Martyn JA, et al. (2005) S-nitrosylation-dependent inactivation of Akt/protein kinase B in insulin resistance. *J Biol Chem* 280: 7511-7518.
83. Pauli JR, Ropelle ER, Cintra DE, Carvalho-Filho MA, Moraes JC, et al. (2008) Acute physical exercise reverses S-nitrosylation of the insulin receptor, insulin receptor substrate 1 and protein kinase B/Akt in diet-induced obese Wistar rats. *J Physiol* 586: 659-671.
84. Neufer PD, Shinebarger MH, Dohm GL (1992) Effect of training and detraining

- on skeletal muscle glucose transporter (GLUT4) content in rats. *Can J Physiol Pharmacol* 70: 1286-1290.
85. Lehnen AM, Leguisamo NM, Pinto GH, Markoski M, De Angelis K, et al. (2011) Exercise-stimulated GLUT4 Expression is Similar in Normotensive and Hypertensive Rats. *Horm Metab Res* 43: 231-235.
 86. Song YJ, Sawamura M, Ikeda K, Igawa S, Nara Y, et al. (1998) Training in swimming reduces blood pressure and increases muscle glucose transport activity as well as GLUT4 contents in stroke-prone spontaneously hypertensive rats. *Appl Human Sci* 17: 275-280.
 87. Chibalin AV, Yu M, Ryder JW, Song XM, Galuska D, et al. (2000) Exercise-induced changes in expression and activity of proteins involved in insulin signal transduction in skeletal muscle: differential effects on insulin-receptor substrates 1 and 2. *Proc Natl Acad Sci U S A* 97: 38-43.
 88. De Angelis Lobo d'Avila K, Gadonski G, Fang J, Dall'Ago P, Albuquerque VL, et al. (1999) Exercise reverses peripheral insulin resistance in trained L-NAME-hypertensive rats. *Hypertension* 34: 768-772.
 89. Milot M, Trudeau F (1997) Plasma galanin immunoreactivity in the rat after swimming. *Physiol Behav* 62: 697-700.
 90. He B, Shi M, Zhang L, Li G, Zhang L, et al. (2011) Beneficial effect of galanin on insulin sensitivity in muscle of type 2 diabetic rats. *Physiol Behav* 103: 284-289.
 91. Thai MV, Guruswamy S, Cao KT, Pessin JE, Olson AL (1998) Myocyte enhancer factor 2 (MEF2)-binding site is required for GLUT4 gene expression in transgenic mice. Regulation of MEF2 DNA binding activity in insulin-deficient diabetes. *J Biol Chem* 273: 14285-14292.
 92. McGee S, Spasling D, Olson A, Hargreaves M (2006) Exercise increases MEF2- and GEF DNA binding activity in human skeletal muscle. *FASEB J* 20: 348-359.
 93. Sparling DP, Griesel BA, Weems J, Olson AL (2008) GLUT4 Enhancer Factor (GEF) interacts with MEF2A and HDAC5 to regulate the GLUT4 Promoter in adipocytes. *J Biol Chem* 283: 7429-7437.
 94. Host HH, Hansen PA, Nolte LA, Chen MM, Holloszy JO (1998) Rapid reversal of adaptive increases in muscle GLUT-4 and glucose transport capacity after training cessation. *84*: 798-802.
 95. Lehnen AM, Leguisamo NM, Pinto GH, Markoski MM, De Angelis K, et al. (2010) The beneficial effects of exercise in rodents are preserved after detraining: a phenomenon unrelated to GLUT4 expression. *Cardiovasc Diabetol* 9: 67.
 96. Mujika I, Padilla S (2000) Detraining: Loss of Training-Induced Physiological and Performance Adaptations. Part I: Short Term Insufficient Training Stimulus. *Sports Medicine* 30: 79-87.
 97. Armstrong RB, Phelps RO (1984) Muscle fiber type composition of the rat hindlimb. *The American Journal of Anatomy* 171: 259-272.
 98. Kump DS, Booth FW (2005) Alterations in insulin receptor signalling in the rat epitrochlearis muscle upon cessation of voluntary exercise. *J Physiol* 562: 829-838.
 99. Reynolds TH 4th, Brozinick JT Jr, Larkin LM, Cushman SW (2000) Transient enhancement of GLUT-4 levels in rat epitrochlearis muscle after exercise training. *J Appl Physiol* 88: 2240-2245.
 100. Mostarda C, Rogow A, Silva IC, De La Fuente RN, Jorge L, et al. (2009) Benefits of exercise training in diabetic rats persist after three weeks of detraining. *Auton Neurosci* 28: 11-16.