

Pseudopelade of BROCQ

Jayakar Thomas*

Department of Dermatology, Chettinad Hospital, and Research Institute, Kelambakkam, Chennai

Corresponding Author*

Jayakar Thomas

Department of Dermatology, Chettinad Hospital,
and Research Institute, Kelambakkam, Chennai

E-mail: jayakarthomas@gmail.com

Copyright: ©2022 Thomas J., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 19 July 2022, Manuscript No. Dmcr-22-17894; **Editor assigned:** 20 July 2022, PreQC No. DMCR-22-17894 (PQ); **Reviewed:** 22 July 2022, DMCR-22-17894, QC No. DMCR-22-17894 (Q); **Revised:** 3 August 2022, Manuscript No. DMCR-22-17894 (R); **Published:** 5 August 2022; Doi: 10.35248/2684-124.22.7.1

Abstract

Pseudopelade of Brocq (PoB) is an unusual form of permanent hair loss from the scalp, the cause of which is unknown. This is a specific diagnosis reached after excluding known causes of patchy scarring alopecia (pseudopelade) such as lichen planopilaris and discoid lupus erythematosus. We present a case of PoB in a 45-year-old female.

Keywords: • Pseudopelade of BROCQ • Patchy • Scarring • Alopecia • Lichen Planopilaris • Discoid lupus • Erythematosus

Introduction

Lowglycaemicrecord

Pseudopelade of Brocq most commonly affects middle-aged and older women. However, it can be present in both sexes and all age groups including children. It has rarely been reported to affect more than one family member. The cause is unknown, as it is the mechanism by which hair loss occurs. Recent studies have shown the cells in the inflammatory infiltrate and the switched-on genes are different in pseudopelade of Brocq compared to lichen Planopilaris and discoid lupus erythematosus, thus providing further evidence this is a distinct entity. Pseudopelade of Brocq is not an inflammatory form of alopecia and the hair loss seems to be due to atrophy of the hair follicles rather than scarring [1].

Clinical features

Pseudopelade of Brocq is often found accidentally as a small patch of hair loss on the scalp. Single hairs may persist in the otherwise hairless patch. The hair loss usually begins at the vertex, the highest point on the scalp. The sides of the scalp (parietal scalp) are another area commonly involved. Rarely has it also been reported in the beard area. See (Figures 1 and 2)

The clinical features of the patches of hair loss are:

- Single or multiple
- Smooth
- Soft
- Slightly depressed
- Round or oval
- Small
- May merge to become larger and irregular in shape.
- Asymptomatic
- Not scaly
- Skin-coloured
- May see pale red/pink around individual hairs initially.
- Hairs may be easily pulled from the edges if active.

The bald areas have been described as looking like 'footprints in the snow.'

The progress of the Pseudopelade of BROCQ is unpredictable however it is generally a slow process. Some patients are left with only one small area of hair loss whereas for others it continues for many years resulting in a significant cosmetic defect. Systemic examination shows no abnormal defects [2].



Figure 1. Footprints in the snow' appearance



Figure 2. Patchy scarring alopecia

Diagnosis

Pseudopelade of Brocq must be distinguished from 'pseudopelade', which is an umbrella term for any form of patchy scarring alopecia such as lichen planopilaris or discoid lupus erythematosus. A scalp biopsy from a smooth patch shows a thin epidermis and sclerotic dermis with streamers of fibrosis down to the fat layer. There is no inflammation seen. Biopsy early in the process may show a small number of lymphocytes around the upper 2/3 of the hair follicle. Braun-Falco proposed clinical and histological criteria for making the diagnosis [3].

Clinical criteria

- Irregular and confluent patches of hair loss
- Moderate atrophy (late-stage)
- Mild redness around hair follicles (early stage)
- Female predominance (3:1)
- A long course of more than 2 years
- Slow progression
- Spontaneous termination possible

Histological criteria

- No marked inflammation
- No widespread scarring
- No significant plugging of hair follicles

- No sebaceous (oil) glands
- Normal epidermis
- Fibrotic streamers in the dermis

Treatment

No treatment has yet been found to stop the process or to regain lost hair. Surgery may help to improve the cosmetic appearance once the condition is no longer active [4-6].

References

1. "Rook Textbook of Dermatology" 7th edn Chapter 63.
2. Yu, M., et al. "Lichen planopilaris and pseudopelade of Brocq involve distinct disease-associated gene expression patterns by microarray". *J Dermatol Sci* (2010): 57: 27-33.
4. Braun.F. O.,et al. "Pseudopelade of Brocq. *Dermatologica*"(1986):172: 18-24.
5. Bitnum, Sam, et al. "Dermatomyositis." *J. pediatr.* 64.1 (1964): 101-131.
6. Flendrie, Marcel, et al. "Dermatological conditions during TNF- α -blocking therapy in patients with rheumatoid arthritis: a prospective study." *Arthritis res. ther.* 7.3 (2005): 1-11.