

## Speciation of Lead and Copper Contents in Non-Alcoholic Wines From Marina Distribution Co. Ltd, Accra, Ghana

Essel Kofi Kwegyir\*

Food Research Institute, Accra, Ghana

### Abstract

The concentrations of two heavy metals (Cu and Pb) were determined in three brands of Ghanaian wines by flame atomic absorption spectrometer (FAAS) after digesting the wine samples with 0.1M HNO<sub>3</sub>. The copper concentration in the three different brands of wine was in the range of 0.4-1.2 mg/L. The toxic lead metal was found to be below the detection (BDL) of 0.1 mg/L. The levels of Cu concentrations were found to be comparable to the permissible limit set by OIV, French and American Wines. The copper and lead identification sources were due to natural sources and anthropogenic sources during vinification processes such as the equipment used.

**Keywords:** Red wine; Resveratrol; *Streptococci*

### Introduction

#### Importance of wine

Wine is one of the most common alcoholic beverages consumed all over the world. Wine naturally contains 85-89% water. Moderate wine consumption can reduce the risk of heart disease and individual components of wine (alcohol and principal polyphenols) can reduce the risk of coronary heart disease in certain individuals. It is important to know that polyphenols can reduce the rate of harmful cell oxidation and favorably affect other processes that, if unchanged, could lead to atherosclerosis and heart disease. Some studies indicated that the cardio protective compounds in grapes, polyphenolic antioxidants, reside in the skin and seeds [1]. Grape skins, which contain purple pigment, are crushed with pulp to make the red wine. The skins give red wine its coloration and contain the highest concentration of polyphenols which are potential antioxidants. Antioxidants are substances that protect cells from oxidative damage caused by free radicals. The polyphenols in red wine includes catechin, gallic acid and epicatechin. Laboratory experiments carried out showed that both wines were active against *streptococci*. The red had a stronger effect than the white, though the difference was not statistically significant. Organic acids in wine, such as acetic acid, citric acid, lactic acid, succinic acid and tartaric acids, are responsible for the antibacterial activity against oral streptococci. The finding suggests that wine “enhances oral health” [2]. Wine polyphenols displayed no activity against oral *Streptococci* or *S. Pyogenes*.

Cell damage caused by free radicals has been implanted in the development of cancer. These chemicals can damage important parts of cells including protein, membranes and DNA. Among other polyphenolic flavonoids, resveratrol is thought to be at least in part responsible for the possible anti-cancer effect of red wine. Resveratrol has been found in at least seventy two plant species, a number of which are components of the human diet such as mulberries, grapes, red wine and peanuts [3-5].

Resveratrol has also shown the ability to inhibit the growth of prostate cancer cells whilst leaving normal cells unaffected [6]. Resveratrol also decreases triglyceride levels as well as low density (LD) “bad” cholesterol levels. Raised levels of serum lipids are strongly associated with an increased risk of cardiovascular disease [7] (Figure 1).

Although, the positive health effects of wine are many, the evidence is clear that these benefits are the highest for those who drink red wine in moderation (two drinks per day for men and one drink per day for women) over extended periods of time. Three or more drinks per day may increase the risk of neurodegeneration, depressive disorders, and obesity, weakening of bones, hypertriglyceridermia, heart disease, hypertension, stroke, breast cancer, suicide and injuries [8]. Wine

consumption in any amount is contraindicated for pregnant women, children and patients with liver disease and in combination with certain medications [9]. Historically, wine has been used as an antiseptic, a pain killer and to treat dermatological conditions and digestive disorders [10].

Regular wine consumption should be used with caution in individuals predisposed to alcoholism, organic diseases and cirrhosis of the liver, migraine, headaches and allergies. Red wine de-alcoholized red wine and grape juice consumption have lowered blood pressure in patients with coronary artery disease or hypertension [11-14].

### Objective

The objective of this technical report is to determine lead and copper contents in white, red and rose wines.

### Literature Review

#### Effects of heavy metals in wine on humans

The identification of heavy metals in wine is a subject of increasing interest since complex atom may reduce their toxicity and bioavailability [15]. Metals in wine can originate from both natural and anthropogenic sources and its concentration can be a significant parameter affecting consumption and conservation of wine. Since metallic ions have important roles in oxido-reductive reactions resulting in wine browning,




Figure 1: Structure of resveratrol.

\*Corresponding author: Kofi Kwegyir Essel, Food Research Institute, Accra, Ghana, Tel: 0285060155/ 0578625912; E-mail: [kofissel@yahoo.com](mailto:kofissel@yahoo.com)

Received August 07, 2014; Accepted November 09, 2014; Published January 03, 2015

Citation: Kwegyir EK (2015) Speciation of Lead and Copper Contents in Non-Alcoholic Wines From Marina Distribution Co. Ltd, Accra, Ghana. Nat Prod Chem Res 3: 157. doi:10.4172/2329-6836.1000157

Copyright: © 2015 Kwegyir EK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

turbidity, cloudiness and astringency, wine quality depends greatly on its metal consumption.

Consumption of wine may contribute to the daily dietary intake of essential metals e.g. copper, iron and zinc, but can also have potentially toxic effects if metal concentrations are not kept under permissible limit [16].

Copper has a definite negative effect on the organoleptic properties of wine [17]. Copper is both an essential and potentially toxic element for humans when in excess. When wine or wine products are consumed in large quantities, the toxic effect of their pollutants may have an additive effect to induction of alcoholism. A typical example is the presence of lead in wines [18]. Several investigators have shown that accumulation of lead in mitochondria of mice occurs with the simultaneous ingestion of small doses of lead and alcohol [19,20]. Trace metal composition of grapes, wines and other alcoholic products is influenced by the type of soil, wine processing equipment vinification methods, the composition of fungicides, insecticides, fertilizers etc. used in the wine industry [21].

Laurie et al. [22] undertook a research titled "Analysis of Major Metallic Elements in Chilean Wines using Absorption Spectroscopy". They concluded that all metal concentrations were within normal ranges as compared to previously reported data from other world wine areas. They also observed that the concentration of Na was higher in the wines produced in the northern parts of the country.

Woldemariam and Chandravanshi [23] studied the concentration of essential and non-essential elements in selected Ethiopian wines. They concluded that wet digestion method and the determination of selected metals at trace levels in wines by flame atomic absorption method were found to be efficient, precise and accurate. It was also observed that Pb concentrations in Ethiopian wines were found in the range of 0.14-0.31 mg/L and these Pb values were somehow larger than the limits set by International Organization for Wines and Grapes (OIV) as 0.2 mg/L. They also said that moderate wine consumption contributes to the daily nutritional requirements of many essential metals. The OIV for Cu is 1 mg/L. The content of some metals can be used to identify the geographic region in which the grapes were grown due to the direct relationship with soil composition [24].

## Method

The wines were analyzed for two heavy metals by AAS and were done in triplicates.

### Analyses by AAS and Flame Photometer

The metals Cu and Pb were determined using Atomic Absorption Spectrophotometer, Perkin Elmer Model 200A using an air/acetylene flame. Distilled water was used to prepare 0.1M HNO<sub>3</sub> acid for dilution of wine solutions. All glasswares were thoroughly washed and were completely dried before use.

### Sampling

For this study, three most popular Ghanaian wine brands consisting of white wine, red wine and rose wines were selected. Seven brands of white wines, six brands of red wines and two brands of rose wines were selected randomly from Marina Distribution Co. Ltd at different sites of Accra, the capital of Ghana. Each bottle of the same three brands was shaken slightly to ensure mixing.

### Ashing

Crucibles were dried in the furnace to a constant weight for 30 minutes. It was crucibles were then cooled to room temperature. The

sample was ashed in a muffle furnace after taking 5.0 ml of the wine samples and evaporation on a hot plate at 550°C for 6-8 hours. The samples in the crucibles were cooled to room temperature after which digestion was carried out.

### Digestion of wine samples

From the bulk samples, 5.0 ml of the wine samples were wet ashed (digested) to decompose the organic substances and make clear solution in duplicates. After digestion was completed, the crucible with the digest was removed from the hot plate and was left to cool (for 10 minutes). The cooled digest was transferred quantitatively into 50 ml volumetric flask after the addition of 0.1 M HNO<sub>3</sub>. The volume of the solution was diluted to the mark with 0.1M HNO<sub>3</sub>. The digest was kept in the refrigerator until the analysis by AAS.

## Results and Discussion

The Lead (Pb) contents in wines found in Marina Distribution Company in Accra were found to be below the detection limit of 0.1 mg/L in all the fourteen wine samples. These lower concentrations of Pb found in the local wines showed that they were of good nutritional value and were of good economic importance if consumed. Pb could be toxic when ingested [25]. Wine is susceptible to lead contamination from the seals on wine bottles and the linings of wine casks.

The concentrations of copper in the studied area ranged between 0.40-1.20 mg/L. This could be compared to the limit set from Marina Distribution Co. Ltd did not contain any lead (Pb) (Table 1).

The analytical wavelengths, the correlation coefficients and correlation equations of the calibration curves for the determination of heavy metals in wine samples by FAAS are given in Table 1. The correlation coefficients of all the calibration curves were > 0.999 and these correlation coefficients showed that there was very good correlation (relationship) between concentration and absorbance. The detection limits for this method for both lead and copper were <0.1 mg/L, which clearly indicate that the method is applicable for the determination of metals at trace levels.

From Table 2, the concentrations of white wines ranged for copper ranged from 0.58-0.89 mg/L with a mean of 0.76 mg/L whilst the red wine concentrations for copper ranged from 0.40-0.84 mg/L with a mean of 0.67 mg/L. The concentration of the Rose wine for copper ranged between 0.50-1.20 mg/L with a mean of 0.85 mg/L. The lead contents in white, red and rose wines were found to be below the detection limit of 0.1 mg/L. The rose wine contains Cu in the highest amounts with concentration 1.2 mg/L. The lowest concentration was found in red wine with copper concentration of 0.40 mg/L.

Metal	Wavelength (nm)	Instrument detection limit	Method detection limit	Correlation coefficient of curve	Equation for calibration curve
Cu	324.7	0.02	0.03	0.9998	Y= 0.0893x - 0.0017
Pb	283.2	0.100	0.10	0.9995	Y= 0.0058x + 0.001

**Table 1:** Analytical wavelengths, detection limits, correlation coefficients and correlation equations of the calibration curves for the determination of metals in wine samples by FAAS. These values were below the standard set by OIV and other countries.

Sample	Cu	Mean	Pb	Mean
White wine	0.58-0.89	0.76	BDL	-
Red wine	0.40-0.84	0.67	BDL	-
Rose wine	0.50-1.20	0.85	BDL	-

**Table 2:** Results of mean concentrations of heavy metals in Ghanaian White, Red and Rose wines (mg/L).

The comparison of heavy metals in Ghanaian wines and published data on wines from different countries is given in Table 3. The Cu contents in Ghanaian wines were in the range of 0.4-1.20 mg/L. The Ghanaian wines had lower levels of Cu than that reported in Czech, Greek, Hungarian, Italian and Spanish wines. However, the Ghanaian Cu contents in wine are comparable to that reported in the French, German and American wines. But the Ghanaian wines contain lower amount of Cu than that reported in Czech, German, Greek, Hungarian, Italian and American wines.

### International Organization for Grapes and Wines (OIV)

Many countries have set maximum permissible limit of some heavy metals in wines considering both the enological and toxicological effects of the heavy metals in wines. The tolerable limits for some heavy metals in wines are also set by International Organization for Grapes and Wines (OIV). The maximum permissible levels in Australia, Germany, and Italy and by OIV are given in Table 4.

The Ghanaian wines contained lower concentration of Cu (0.4-1.2 mg/L) than the limit set for Cu in Australia, Germany and Italy. However, the Ghanaian content of Cu in wine is comparable to that set for Cu (1 mg/L) by OIV. The Pb contents were below the detection limit in wines from Ghanaian market and cannot be compared to the permissible limits set by OIV and the countries in Table 4.

Metal	Concentrations of heavy metals (mg/L) in different countries	
Country/Metal	Copper (Cu)	Lead (Pb)
Czech	0.012-6.827	0.010-1.253
French	ND-0.48	0.006-0.023
German	0.02-0.71	-
Greek	0.2-1.65	ND-0.62
Hungarian	0.15-2.57	-
Italian	0.001-1.34	0.01-0.35
Spanish	ND-3.1	0.001-0.096
American	0.05-0.58	-
Ghanaian	0.4-1.2	BDL

**Table 3:** Comparison of the concentrations of heavy metals in wines of different countries.

Country	Concentration of metals (mg/L)				
	Na	Cu	Zn	Pb	Cd
Australia		5	5	0.2	0.05
Germany		5	5	0.3	0.01
Italy		10	5	0.3	0.01
OIV	0.4-1.2	BDL of 0.1 mg/L			

**Table 4:** Comparison of the levels of metals in Ghanaian wines to permissible levels of some metals (mg/L) in some countries and by OIV.

### Conclusion

The Ghanaian wines were found to contain copper values lower than the limits set by OIV. The concentrations of white and red wines for copper ranged from 0.58-0.89 mg/L with a mean value of 0.76 mg/L and the red wine for copper with a range of 0.40- 0.84 mg/L with a mean value of 0.67 mg/L Except Czechoslovakia, which had a range higher than that set by OIV (0.012- 6.827 mg/L). This shows that wines could be a good source of Cu for humans. The toxic Lead was not detected in Ghanaian wines revealing that Ghanaian wines contain very low concentration of Pb (<0.1 mg/L). In general the Cu content values in wine are in line with the previously reported data set by OIV, American and French wines. The Ghanaian wines are safe for consumption and also good for economic reasons.

### References

- Gabriela V, Gabriel D, Radoi B (2011) Determination of antioxidant capacity for some products based on wine and medicinal plants. Journal of Agroalimentary processes and Technologies 17: 151-156.
- Daglia M, Papetti A, Grisoli P, Aceti C, Dacarro C, et al. (2007) Antibacterial activity of red and white wine against oral streptococci. J Agric Food Chem 55: 5038-5042.
- Langcake P, Pryce RJ (1976) The production of resveratrol by *Vitis vinifera* and other members of the vitaceae as a response to infection or injury. Physiol Plant Pathol 9: 77-86.
- Sanders TH, McMichael RW Jr, Hendrix KW (2000) Occurrence of resveratrol in edible peanuts. J Agric Food Chem 48: 1243-1246.
- Slemani EH, Creasy LL (1992) Concentration of Phytoalexin resveratrol in wine. Am J Enol Vitic 43: 49-52.
- Narayanan NK, Narayanan BA, Nixon DW (2004) Resveratrol-induced cell growth inhibition and apoptosis is associated with the modulation of phosphoglycerate mutase B in human prostrate cancer cells: two dimensional sodium dodecyl sulphate- polyacrylamide gel electrophoresis and mass spectrometry evaluation. Cancer Detect Prev 28: 443-452.
- Miura D, Miura Y, Yagasaki K (2003) Hypolipidemic action of dietary resveratrol, a phytoalexin in grapes and red wine, in hepatoma-bearing rats. Life Sci 73: 1393-1400.
- Saremi A, Arora R (2008) The cardiovascular implications of alcohol and red wine. Am J Ther 15: 265-277.
- Fehér J, Lengyel G, Lugasi A (2005) [Cultural history of wine, the theoretical background of wine therapy]. Orv Hetil 146: 2635-2639.
- Robinson J (2006) The Oxford companion to wine. Oxford University Press, New York.
- Foppa M, Fuchs FD, Preissler L, Andrighetto A, Rosito GA, et al. (2002) Red wine with the noon meal lowers post-meal blood pressure: a randomized trial in centrally obese, hypertensive patients. J Stud Alcohol 63: 247-251.
- Park YK, Kim JS, Kang MH (2004) Concord grape juice supplementation reduces blood pressure in Korean hypertensive men: double-blind, placebo controlled intervention trial. Biofactors 22: 145-147.
- Karatzis KN, Papamichael CM, Karatzis EN, Papaioannou TG, Aznaouridis KA, et al. (2005) Red wine acutely induces favorable effects on wave reflections and central pressures in coronary artery disease patients. Am J Hypertens 18: 1161-1167.
- Jiménez JP, Serrano J, Taberner M, Arranz S, Díaz-Rubio ME, et al. (2008) Effects of grape antioxidant dietary fiber in cardiovascular disease risk factors. Nutrition 24: 646-653.
- Cocchi M, Franchini G, Manzini D, Manfredini M, Marchetti A, et al. (2004) A chemometric approach to the comparison of different sample treatments for metals determination by atomic absorption spectroscopy in aceto Balsamico tradizionale di Modena. J Agric Food Chem 52: 4047-4056.
- Tariba B (2011) Metals in wine—impact on wine quality and health outcomes. Biol Trace Elem Res 144: 143-156.
- Gennaro MC, Mentasti E, Sarzanini C, Pestico A (1996) Undesirable and harmful metals in wines, determination and removal. Food Chem 19: 93-104.
- Roses OE, Gonzalez DE, Lopez CM, Pinerio AE, Villamil EC (1997) Lead levels in argentine market wines. Bull Environ Contam Toxicol 59: 210-215.
- Brivet I, Chomard PH, Dumas P, Lallemand AM, Theven M (1990) Ethanol-lead in the Sprague Dawley rat. Food Addit Contam 71: S150-S151.
- Nation JR, Dagger LM, Dwyer KK, Braston GR, Grorer CA (1999) The effects of dietary lead in ethanol-reinforced responding. Alcohol-Alcoholism 26: 473-480.
- Eschauer H, Neeb RE (1988) Micro-element analysis in wine and grapes. Modern Methods of Plant Analysis 6: 67-91.
- Felipe Laurie V, Evelyn Villagra, Jamie Tapia, Jorge Sarkis ES, Marcos AH (2010) Analysis of major metallic elements in Chilean wines by Atomic spectroscopy. Gen Inv Agr 37: 77-85.
- Daniel M, Woldemariam B, Chandravanshi S (2011) Concentration levels of Essential and Non-essential Elements in selected Ethiopian wines. Chemical Society of Ethiopia 25: 168-180.
- Pryczynska K (2007) Chemical speciation and fractionation of metals in wine. Chem Spec Bioav 19: 1-8
- Lara R, Cerutti S, Salonia JA, Olsina RA, Martinez LD (2005) Trace element determination of Argentine wines using ETAAS and USN-ICP-OES. Food Chem Toxicol 43: 293-297.